



Cervid Solutions-Vaccination Schedule

Spring:

New to vaccination program

- Bred does
 - 2cc - 50-60 days prior to fawning
 - 2cc – Booster, 2-4 weeks after dose 1
- Males:
 - 2cc - Anytime between February 15st to April 1st
 - 2cc - Booster, 2-4 weeks after first dose

Already on the Cervid Solutions program

- Bred does
 - 2cc -30 days prior to fawning
- Males
 - 2cc – Anytime between February 15st to April 1st

See bottle-fed fawn protocol for information on fawn vaccination schedule

Fall:

New to Program

- Females:
 - 2cc - Upon weaning of fawns or at CIDR insertion
 - 2cc - Booster 2-4 weeks after first dose
- Males:
 - 2cc- After antler growing season and prior to breeding
 - 2cc - Booster 2-4 weeks after first dose
- Fawns
 - Pre-Weaning Article link <http://www.cervidsolutions.com/961>
 - 1cc - 3 weeks prior to weaning
 - 1cc= Booster upon weaning (Physical Separation from Mom)

Already on program

- Females
 - 2cc - Upon fawn weaning/separation for breeding or CIDR insertion for A.I.
- Males:
 - 2cc - After antler growing season and prior to breeding (September/October)

Recap-

All animals, when they begin the program, need a booster dose 2-4 weeks after first dose. After they have started the program and with initial, plus booster dose, they transition to the maintenance protocol, delivering 2cc bi-annually with one dose in Fall and one dose in Spring.